

THE BELGIAN VIRTUAL T U M O U R B A N K

Newsletter / December 2020

In March 2008, the nationwide virtual tumourbank project was launched by the Belgian Minister of Health, Ms. Laurette Onkelinx (initiative 27 of the Belgian Cancer Plan). The Belgian Virtual Tumourbank (BVT network) encompasses the tumour biobanks from eleven Belgian university hospitals (<u>click here</u> for the list) that collect and store residual human tumour samples locally. In order to facilitate the search for tumour samples scattered among different Belgian institutions, data collected at sample level is made available for researchers via the online BVT catalogue (BVTc) application. A high quality of the data on the tumour samples requested by scientists for research in oncology is guaranteed by automatic and manual controls performed by the BVT project team at the Belgian Cancer Registry.


METASTASIS SAMPLES IN THE BVT CATALOGUE

In 2018 about one in five cancers in Belgium for which the TNM classification is applicable, were already metastasised on the moment of diagnosis (cM1 or pM)¹. The BVT catalogue can facilitate the search for tumour samples scattered among different local biobanks, especially for less common sample types such as metastasis samples. If you want to know if the specific samples for your research are available in our catalogue, please request <u>full access</u> so you can browse freely through all available sample data and locate the matching samples for your study, or fill in a <u>Sample Availability Request form</u> on our website!

WHICH METASTASIS SAMPLES ARE AVAILABLE?

In total, 107,935 registrations were available in the BVT catalogue on the moment of selection (October 2020), of which **13,235** registrations from metastasis samples from **6,814** patients. The BVT catalogue contains slightly more registrations from metastasis samples that originate from male patients (50.8%) than from female patients (49.2%).

The most common sample localisations of the metastasis samples in the BVT catalogue are liver (26.0%) and lymph nodes (25.8%). Soft tissue completes the top three of sample localisations with 10.9%, followed by central nervous system (10.5%) and lung (9.3%) (see figure 1).





FROM WHICH PRIMARY TUMOUR LOCALISATIONS DO THE METASTASES ORIGINATE?

The metastasis samples most commonly originate from primary tumours of colon and rectum (18.5%), lung (7.9%), breast (7.8%) and skin (7.4%) (see figure 2). For 34.6% of the patients the localisation of the corresponding primary tumour is unknown.




Figure 2: Overview of the most common primary tumour localisations of metastases in the BVTc

For approximately 10% of the patients with metastasis samples registered in the BVTc also a corresponding primary tumour tissue sample is available.


WHICH MATERIAL TYPES ARE STORED BESIDES TISSUE OF THE METASTASIS?

For 68.9% of the registrations, only residual tumour tissue is stored. For some patients, also additional types of material are stored at the local biobanks and registered in the BVT catalogue. The most common types are corresponding normal tissue (21.7%) and blood (9.2%). Plasma (6.8%), serum (5.2%), buffy coat (1.8%) and DNA (1.6%) are also available in some local biobanks.

As far as the conservation is concerned, 74.1% of the metastasis samples is stored at -80°C, while 25.9% is included in paraffin blocks.

SPREAD THE WORD....

Over the past year, the Belgian Virtual Tumourbank team presented a poster at the BACR annual meeting (7th February 2020, Brussels) and two posters at the Europe Biobank Week (17-20th November 2020, online).



BIOBANK WEEK 2020 17 El Number Votal Calenares Biologiani, cirál 2 Rents Ultimate	Belgian Virtual Tumourbank (BVT Additional data quality control kidney cancer samples registered Both, the samples and the samples government in a bage with Theoretail government in a bage with Theoretail government in a bage with the samples monotone and the samples and the samples monotone and the samples and the samples monotone and the samples	ON In BVT estate on behalf of	-1	44
INTRODUCTION	AIM	METHODS		
The below Workel Termshood XMM (In steel termshood terms terms standing to redsh termshood terms terms terms termshood termshood termshood terms	Investigate and improve the data quality of kickey cancer samples from sample years 2017 and 2016 stored in the patalogue of the Brights Virtual Turneurbark by performing an additional data quality control	The BVT analysis data of all storage cancer cannot force 2012-2013 to compared to the data multiple data the CD server registry distance. • Unlogar based on the solid storage high control registry distance. • Unlogar based on the solid security identification running (SSM) is the solid storage of		
	RESULTS			
	In September 2020, a total of 207,657 registrations were available in the 50/11 of which 664 registrations from primary kidney canon samples from the years 2017 2020. The latter argents from 100 kidney canon cases argent to 10 our of 11 biobents of the 507 motions.	Variables	Cancer cases registered in BVT = (%)	Corrections needed in BVT in (%)
		SSIV	563 [330]	D
	Cut of the 12 services that seems takes that access the this access the three of the three one of the terror three in the terror three integration, different takes periods on the terror three terror three terrors the terror three terrors the terror tere	Birth-dute Georger	265 [200]	0
		Seeder Surriedate	263 (100)	10.5
		Bepty number	263 (100)	112.19
		Sartaly form	263 (200)	
		Sample localization	263 (200)	10.9
		Histories	263 (200)	15/5/0
		Balaning.	263 (100)	100
gh quality of the data is guaranteen by outpenatic and manual controls	registered renal turneer turned out to be non-involve (behaviour "2").	Labourty	245 (25.5)	50.20
QC1: automatic check for format and content by the BVI		Differentiation ande	141(05.5)	50.0
application GC2: manual check for inconsistencies between different fields by	database showed that	aT.	236-88.73	70.0
BVT data analysis, but these initial contrain, the data is published in the BVT tatalogue of made available for measurbers in a coded form.	 two patients recorned neoadjacant the rapy prior to sampling 	Additional information		added in the BVT
and made available for researchers in a coded form. 1. OC1: only performed for a sanoffic data quality control mudy or upon		Recurrent turnous by TNMI Turn our resection offer neoadjacant therapy (suTNM)		50.99
(g) a period personnel for a specific cash spacify control study or upon sparst from a resolution on state of specific samples solution for spheri study.				2 (0.0)
CONCLUSIONS	AKNOWLEDGEMENTS	CONTACT INFORMATION		
The WT darkson with fund to have a generative of data. Since this much spectrum alternative sector on course of the data sector and matchenet. Denays piece to reservice of the knew taneous have fund to make them to ensure the altern evolution of the base and course ingetter referentiation and parameters a high quality of sourceast data from the presence of the transition of a source transition.		waran di unterpretanti be Region Vitual Tanoarbani - Segion Coster Regiony Resegion 21 Sector 2015 tes 7 32 di Branch Indeed Reservation og		

Flyers on the BVT project have been distributed in 2020 at the BACR annual meeting (7th February 2020, Brussels), the BSMO annual meeting (14-15th February, Ghent) and the Belgian Week of Gastroenterology (4-6th March, Antwerp).

Please let us know if you are aware of upcoming meetings where the BVT project can be presented or if you would like to receive BVT flyers for distribution in your institution.

For information on sample availability in the 11 biobanks of the BVT network, click here.

To gain full access to the online BVT catalogue, please follow the procedure indicated here.

For further information: <u>www.virtualtumourbank.be</u> <u>biobank@kankerregister.org</u> or <u>biobank@registreducancer.org</u>

Do you wish to be kept informed in the future? Do you wish not to receive our information any longer? Send an email to <u>biobank@kankerregister.org</u> Belgian Cancer Registry | Brussels | <u>www.kankerregister.org</u>